




A service of the U.S. National Library of Medicine  
and the National Institutes of HealthMy NCBI   
[\[Sign In\]](#) [\[Register\]](#)All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books  
Search PubMed  for   [Advanced Search](#)[Limits](#) [Preview/Index](#) [History](#) [Clipboard](#) [Details](#)Display [AbstractPlus](#)  Show [20](#)  Sort By  Send to [All: 1](#) [Review: 0](#) 1: AIDS. 1997 Dec;11(15):1799-806.  [Links](#)**AIDS prognosis based on HIV-1 RNA, CD4+ T-cell count and function: markers with reciprocal predictive value over time after seroconversion.****de Wolf F, Spijkerman I, Schellekens PT, Langendam M, Kulken C, Bakker M, Roos M, Coutinho R, Miedema F, Goudsmit J.**

Department of Human Retrovirology, University of Amsterdam, The Netherlands.

**OBJECTIVE:** HIV-1 RNA levels in peripheral blood are strongly associated with progression to AIDS, CD4+ T-cell decline, or death. Their predictive value is reportedly independent of the predictive value of CD4+ T-cell counts. Because the interrelations between these parameters of HIV-1 infection are poorly understood, we studied the kinetics and predictive value of serum HIV-1 RNA levels, CD4+ T-cell counts, and T-cell function. **DESIGN AND METHODS:** HIV RNA levels, CD4+ T-cell counts, and T-cell function were measured from seroconversion to AIDS in 123 homosexual men who seroconverted during a prospective study and were followed over 10 years. **RESULTS:** Two patterns of median HIV-1 RNA levels were found during infection: a steady-state and a 'U-shaped' curve. Steady-state high RNA levels were related to rapid disease progression. For the U-shaped curve, there were groups with high and low RNA levels related to disease progression. At 1 year after seroconversion, RNA level was the only marker that was strongly predictive. Furthermore, decreasing RNA levels in the first year following seroconversion were related to better prognosis than stable low levels. Low CD4+ T-cell count and T-cell function became predictive of progression to AIDS at 2 and 5 years after seroconversion, respectively. **CONCLUSIONS:** With ongoing infection, the predictive value of low CD4+ T-cell count and T-cell function increases, whereas the predictive value of high HIV-1 RNA level decreases. These findings reflect the observation that infection with HIV progressively leads towards immune deficiency, which in later stages is most predictive of disease progression.

PMID: 9412697 [PubMed - indexed for MEDLINE]

**Related Articles**

Early and late HIV-1 RNA level and its association with other markers and disease progression in long-term AIDS-free homosexual men. [AIDS. 1997]

Persistent immune activation in HIV-1 infection is associated with progression to AIDS. [AIDS. 2003]

CD4+ and CD8+ lymphocytes and HIV RNA in HIV infection: high baseline counts and in particular rapid decrease of CD8+ lymphocytes predict AIDS. [AIDS. 1999]

Review Immunologic and virologic markers determining progression to AIDS. [Acquir Immune Defic Syndr Hum Retroviro. 1995]

Review The role of viral diversity in HIV pathogenesis. [Acquir Immune Defic Syndr Hum Retroviro. 1995]

[» See Reviews...](#) | [» See All...](#)**Recent Activity**[Turn Off](#) [Clear](#)

AIDS prognosis based on HIV-1 RNA, CD4+ T-cell count and function: markers with reciprocal...

Prognostic value of HIV-1 RNA, CD4 cell count, and CD4 Cell count slope for progression to...

[CD4 count AND HIV infecti... \(406\)](#)

Association Between HIV-1 RNA Level and CD4 Cell Count Among Untreated HIV-Infected Indivi...

[CD4 count AND viral load... \(103\)](#)

PubMed

Display [AbstractPlus](#)  Show [20](#)  Sort By  Send to [Write to the Help Desk](#)[NCBI](#) | [NLM](#) | [NIH](#)[Department of Health & Human Services](#)[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)